Natural Resources Canada | Ressources naturelles Canada

GEOLOGICAL SURVEY OF CANADA
OPEN FILE 8370

Triassic formations in Liard Basin, northeastern British Columbia: sample descriptions, and detrital zircon U-Pb geochronology, methodology, results, and data tables

M.E. McMechan, F. Ferri, and W. Matthews

2018

Celebrating 175 yrs

Canada

GEOLOGICAL SURVEY OF CANADA
OPEN FILE 8370

Triassic formations in Liard Basin, northeastern British Columbia: sample descriptions, and detrital zircon U-Pb geochronology, methodology, results, and data tables

M.E. McMechan[1], F. Ferri[2], and W. Matthews[3]

[1] Geological Survey of Canada, 3303-33rd Street NW Calgary, Alberta T2L 2A7
[2] Petroleum Geoscience Section, Tenure and Geoscience Branch, Ministry of Natural Gas Development, Victoria, British Columbia V9A 5N7
[3] Centre for Pure and Applied Tectonics and Thermochronology, Department of Geosciences, University of Calgary, 2500 University Drive NW, Calgary, Alberta T2N 1N4

2018

© Her Majesty the Queen in Right of Canada, as represented by the Minister of Natural Resources, 2018

Information contained in this publication or product may be reproduced, in part or in whole, and by any means, for personal or public non-commercial purposes, without charge or further permission, unless otherwise specified.
You are asked to:
- exercise due diligence in ensuring the accuracy of the materials reproduced;
- indicate the complete title of the materials reproduced, and the name of the author organization; and
- indicate that the reproduction is a copy of an official work that is published by Natural Resources Canada (NRCan) and that the reproduction has not been produced in affiliation with, or with the endorsement of, NRCan.

Commercial reproduction and distribution is prohibited except with written permission from NRCan. For more information, contact NRCan at nrcan.copyrightdroitdauteur.rncan@canada.ca.

Permanent link: https://doi.org/10.4095/306629

This publication is available for free download through GEOSCAN (http://geoscan.nrcan.gc.ca/).

Recommended citation
McMechan, M.E., Ferri, F., and Matthews, W., 2018. Triassic formations in Liard Basin, northeastern British Columbia: sample descriptions, and detrital zircon U-Pb geochronology, methodology, results, and data tables; Geological Survey of Canada, Open File 8370, 1 .zip file. https://doi.org/10.4095/306629

Publications in this series have not been edited; they are released as submitted by the author.

ABSTRACT 2
INTRODUCTION 3
TRIASSIC SAMPLE INFORMATION 5
Ludington Formation 5
Liard Formation 7
Toad Formation 8
Grayling Formation 9
DETRITAL ZIRCON U-PB AGE DETERMINATION METHODS 11
ACKNOWLEDGEMENTS 12
REFERENCES 13

ABSTRACT

In the Liard Basin of western Canada, Lower and Middle Triassic aged strata comprise a northeasterly sourced, coarsening upward clastic wedge that is overlain by mixed clastic and carbonate units of Late Ladinian to Norian age. This report provides sample information, detrital zircon U-Pb geochronology methodology, results and data tables for a suite of seven samples collected from Lower Triassic to Carnian formations in western Liard Basin. These data show there was no western sourced material and suggest that the clastic material was recycled primarily from the Caledonian and Ellsemerian clastic wedges and the high Arctic. When combined with previously published results for 5 additional samples, they provide a robust reference frame for the Lower Triassic to Carnian succession of the Liard basin.

INTRODUCTION

The Liard Basin, a sub-basin of the Western Canada Sedimentary Basin, is characterized by anomalously thick Upper Paleozoic and mid-Cretaceous strata (Leckie et al., 1991; Wright et al., 1994). The Basin, bounded on the east by the deep-seated Bovie Lake Structure (Figure 1), contains up to 5 km of Phanerozoic strata overlying Proterozoic strata and the Laurentian craton. In Liard Basin the Triassic succession thins markedly from a maximum thickness of approximately 1600 m in the west to a couple of hundred metres in the northeast. Thinning is primarily due to increased erosion beneath the sub-Cretaceous unconformity. Lower and Middle Triassic strata form a northeasterly sourced clastic succession that changes up section from siltstone and shale-dominated (Grayling Formation), to interlayered siltstone, shale and sandstone (Toad Formation) to sandstone dominated (Liard Formation). Younger Middle Triassic and Upper Triassic (late Ladinian to Norian) strata are mixed clastic and carbonate (Ludington and Pardonet formations) and carbonate (Baldonnel Formation) units. These are only preserved beneath the sub-Cretaceous unconformity in western exposures (Figure 2). To a large extent the Grayling and Toad formations are facies equivalents (Gibson et al., 1991). The Liard Formation changes facies westward into the Ludington and Toad formations (Taylor and Stott, 1973; Gordey et al., 1992). McMechan et al. (2012) provide a more detailed description of Triassic Formations exposed in the British Columbia part of the Liard Basin.



Figure 1. Location of Triassic aged detrital zircon samples analysed.

Seven samples were collected from Triassic strata as part of a larger study examining the detrital zircon provenance of the Liard Basin (McMechan et al., 2015, 2017). These samples are from a broad

stratigraphic interval and included some from the westernmost exposures of the Liard Basin (Figure 1, 2). They were collected to determine: 1) if there was a significant change in detrital zircon U-Pb age signature with stratigraphic level; and 2) if there was a component of west-derived Triassic or upper Paleozoic (< 350 Ma) detrital zircon grains, as had been reported by Beranek and Mortenson (2011) for uppermost Middle Triassic (late Ladinian) ancestral North American margin strata in southeast Yukon 300 km to the northwest. Unlike the Ladinian (uppermost Middle Triassic) Laurentian continental margin strata sampled by Beranek and Mortenson (2011) our samples contained only one younger (< 360 Ma) western sourced grain. Our results show little variation with stratigraphic level. They also indicate recycling of material derived from the Caledonian and Ellsemerian clastic wedges and high Arctic (Figure 3). When combined with published results from 4 samples of the Toad Formation (Golding et al., 2016) and from one sample from the Liard Formation (Gehrels and Pecha, 2014) they provide a robust reference frame for the Lower Triassic to Carnian succession of the Liard basin.



Figure 2. Schematic stratigraphic cross section of Triassic strata across the southern Liard Basin. Red stars show approximate stratigraphic position of detrital zircon samples. See Figure 1 for location.



Figure 3. Probability density plot for the combined data set for all samples of Triassic strata in this study.

TRIASSIC SAMPLE INFORMATION

The age estimate provided for each sample is based on Figure 8.55 by E.T. Tozer in Gordey et al. (1992).

Ludington Formation

Detrital zircon U-Pb ages were obtained for two samples of the Ludington Formation from exposures in the western part of the basin (Figure 1, 2). Sample 14MBB001B01 (Figure 4) came from the base of the formation near the southwestern end of the basin and sample 14MBB026B01 (Figure 5) came from 50 m below the Cretaceous unconformity in the central part of the basin.

Field No. 14MBB001B01; GSC Curation No. C-591547; CAPPT lab number: UK021

Lithology: medium brown grey weathering, dark grey, laminated, calcareous, very coarse grained siltstone.

Location: 0365335E, 6525600N NAD83, Zone 10

Estimated age: Carnian

Figure 4. Ludington Formation outcrop sampled for sample 14MBB001B01.

Field No. 14MBB026B01; GSC Curation No. C-591540; CPATT lab number: UK019

Lithology: light grey to brown grey weathering, medium grey, calcareous, very fine grained, quartzose sandstone.

Location: 0363526E, 6560859N NAD83, Zone 10.

Estimated age: Carnian



Figure 5. Ludington Formation outcrop sampled for sample 14MBB026B01.

Liard Formation

Detrital zircon U-Pb ages were obtained for one sample of the Liard Formation from the basal part of the formation along Toad River (Figure 1, 2, 6).

Field No. 13EPF008B01; GSC Curation No. C-571528; CPATT lab number: UK012

Lithology: buff weathering, dark grey, silty, calcareous fine to very fine grained sandstone in medium to thin beds. Character of the Liard Formation along Toad River shown in Figure 6.

Location: 0402654E, 6561908N NAD83, Zone 10

Estimated age: Ladinian

Figure 6. Liard Formation exposed across Toad River from the outcrop sampled for sample 13EPF008B01.

Toad Formation

Detrital zircon U-Pb ages were obtained for two samples of the Toad Formation from exposures in the western part of the basin (Figure 1, 2). Both samples came from near the base of the formation along the Alaska Highway. Sample 14MBB005C01 (Figure 7) came from the first Triassic exposures west of Toad River Lodge and sample 14MBB003A01 (Figure 8) came from the westernmost exposure of the Liard formation south of Liard River Hot Springs.

Field No. 14MBB005C01; GSC Curation No. C-591550; CPATT lab number: UK005

Lithology: brown weathering, grey brown, very fine grained silty sandstone with some calcite veins.

Location: 04366196E, 6525918N NAD83, Zone 10

Estimated age: Nammalian

Figure 7. Sandstone at base of the Toad Formation on the Alaska Highway west of Toad River Lodge sampled for sample 14MBB005C01.

Field No. 14MBB003A01; GSC Curation No. C-591554; CPATT lab number: UK006

Lithology: tan brown weathering, dark grey, calcareous silty, very fine grained sandstone to sandy siltstone, generally laminated with some thin 2 to 5 cm thick siltier bioturbated layers.

Location: 031197E, 6585714N NAD83, Zone 10

Estimated age: Nammali

Figure 8. Sandstones near the base of the Toad Formation on the Alaska Highway south of Liard Hot Springs. Sample 14MBB003A01came from the bed behind the person.

Grayling Formation

Two samples of the sandstone from within 3 m of the base of the Grayling Formation (Figure 9, 10) were analysed from exposures in the western part of the basin (Figure 1, 2).

Field No. 14MBB027B01; GSC Curation No. C-591542; CAPPT lab number: UK015

Lithology: light brown weathering, light grey, very fine grained sand with the occasional mica flake; very thick to thick bedded locally with low angle lamination.

Location: 04366196E, 6525918N NAD83, Zone 10

Estimated age: Griesbachian

Figure 9. Sandstone at base of Grayling Formation sampled by sample 14MBB027B01.

Field No. 14MBB029A01; GSC Curation No. C-591544; CPATT lab number: UK020

Grey brown weathering, light grey, very fine grained sandstone, medium bedded, barely calcareous, weathers to rubble.

Location: 0355903E, 6603832N NAD83, Zone 10

Estimated age: Griesbachian

Figure 10. Sandstones at base of Grayling Formation. Sample 14MBB029A01came the rubbly sandstone intervals to the right and left of the sample bag.

DETRITAL ZIRCON U-PB AGE DETERMINATION METHODS

U-Pb isotopic data was collected by LA-ICP-MS methods. Seven samples were processed and measured at the Centre for Pure and Applied Tectonics and Thermochronology (CPATT), University of Calgary, Calgary, Canada (full details of the measurement method are presented in Matthews and Guest (2016)). Dates and probablity of concordance were calculated for each analysis using the Excel™ plug-in Isoplot v. 4.15 (Ludwig 2012). Measurements with a probability of concordance of less than 1% were eliminated from the dataset. The distribution of the 586 dates that met our filtering criteria are presented on a probability density plot (Figure 3).

ACKNOWLEDGEMENTS

We thank Christine Deblonde for assistance in the field, Jayden Tse and Dave Sargent for technical assistance and Lisel Currie for her review of this open file. This study was funded by the Geological Survey of Canada. Detrital zircon analyses were conducted at the Centre for Pure and Applied Tectonics and Thermochronology at the University of Calgary in an laser ablation–inductively coupled plasma–mass spectrometry laboratory that is funded by the Canadian Foundation for Innovation (CFI project 30696).

REFERENCES

Beranek, L.P., and Mortensen, J.K., 2011. The timing and provenance record of the Late Permian Klondike orogeny in northwestern Canada and arc-continent collision along western North America. Tectonics, vol. 30, TC5017. doi:10.1029/2010TC002849.

Gehrels, G., and Pecha, M., 2014. Detrital zircon U-Pb geochronology and Hf isotope geochemistry of Paleozoic and Triassic passive margin strata of western North America; Geosphere, vol. 10, p. 49-65. doi:10.1130/GES00889.

Golding, M.I., Mortensen, J.K., Ferri, F., Zonneveld, J.-P., and Orchard, M.J., 2016. Determining the provenance of Triassic sedimentary rocks in northeastern British Columbia and western Alberta using detrital zircon geochronology, with implications for regional Tectonics; Canadian Journal of Earth Sciences, vol. 53, p. 1-16.

Gordey, S.P., Geldsetzer, H.H.J., Morrow, D.W., Bamber, E.W., Henderson, C.M., Richards, B.C., et al., 1992. Upper Devonian to Middle Jurassic assemblages, Part. A, Ancestral North America; *in* Geology of the Cordilleran Orogen in Canada. Gabrielse, H., and Yorath, C.J. (Ed.); Geological Survey of Canada, Geology of Canada no. 4, p 221–328.

Leckie, D.A., Potocki, D.J. and Visser, K. 1991. The lower cretaceous Chinkeh formation: A frontier-type play in the Liard basin of western Canada; American Association of Petroleum Geologists, Bulletin 75, p. 1324-1352.

Ludwig, K.R., 2012, Isoplot v. 3.75—a geochronological toolkit for Microsoft Excel: Berkeley Geochronology Center, Special Publication, no. 5, p. 1-75.

Matthews, W.A., and Guest, B. 2017. A practical approach for collecting large n detrital zircon U-Pb datasets by quadrupole LA-ICP-MS; Geostandards and Geoanalytical Research, vol. 41, p. 161-180. doi: 10.1111/ggr.12146.

McMechan, M., Ferri, F. and MacDonald, L. 2012. Geology of the Toad River area (NTS 094N), northeast British Columbia; *In* Geoscience Reports 2012. British Columbia Ministry of Energy and Mines, pp. 17–39.

McMechan, M., Currie, L., Matthews, W., Ferri, F. and Guest, B., 2015. Detrital zircon geochronology and the evolution of clastic source areas, Liard Basin, northeast British Columbia; Geoconvention 2015, abstract.
http://www.geoconvention.com/archives/2015/250_GC2015_Detrital_zircon_geochronology.pdf [accessed April 11, 2016].

McMechan, M., Currie, L., Ferri, F., Matthews, W. and O'Sullivan, P. 2017. Cambrian detrital zircon signatures of the northern Cordilleran passive margin, Liard area,Canada: evidence of sediment recycling, non-Laurentian ultimate sources, and basement denudation; Canadian Journal of Earth Sicences, v. 54, p. 609-621.

Taylor, G.C. and Stott, D.F. (1973): Tuchodi Lakes map-area, British Columbia; Geological Survey of Canada, Memoir 373, 37 p .

Wright, G.N., McMechan, M.E., Potter, D.E.G. and Holter, M.E. 1994. Chapter 3 - Structure and Architecture of the Western Canada Sedimentary Basin; *in* Geological Atlas of the Western Canada Sedimentary Basin, Mossop, G.D. and Shetsen, I. (comp.). Canadian Society of Petroleum Geologists and Alberta Research Council, p. 25-40.